

DUAL INHIBITION OF POST-RADIOGENIC ANGIO- VASCULOGENESIS BY OLAPTESED PEGOL (NOX-A12) AND BEVACIZUMAB IN GLIOBLASTOMA – INTERIM DATA FROM THE FIRST EXPANSION ARM OF THE GERMAN PHASE 1/2 GLORIA TRIAL. Frank A. Giordano¹, Julian P. Layer^{1,2}, Sonia Leonardelli², Lea L. Friker³, Christina Schaub⁴, Roberta Turiello², Elena Sperk⁵, Iris Mildenberger⁶, Franziska Grau⁷, Daniel

¹ Department of Radiation Oncology, University Hospital Bonn; ² Institute of Experimental Oncology, University Hospital Bonn; ³ Department of Neurology, University Hospital Bonn; ³ Department of Radiation Oncology, University Hospital Bonn; ⁴ Department of Neurology, University Hospital Bonn; ⁵ Department of Radiation Oncology, University Hospital Bonn; ⁴ Department of Neurology, University Hospital Bonn; ⁵ Department of Radiation Oncology, University Hospital Bonn; ⁴ Department of Neurology, University Hospital Bonn; ⁵ Department of Radiation Oncology, University Hospital Bonn; ⁴ Department of Neurology, University Hospital Bonn; ⁴ Department Abbaa, ⁴ Department of Neurology, University Hos Mannheim, University of Heidelberg; ⁶ Department of Neurology, University Hospital Mannheim, University Hospital Bonn; ⁸ Institute of Neuropathology, University Hospital Leipzig; ⁹ Department of Neurology, University Hospital Münster; ¹⁰ Department of Neurology, University Hospital Tübingen; ¹¹ Department of Neurology, University Hospital Essen; ¹² Department of Neuroradiology, National Hospital for Neurology London; ¹³ Department of Radiotherapy, University Hospital Leipzig



Copies of this poster are for personal use only and may not be reproduced without permission from the authors of this post

Paech⁷, Torsten Pietsch³, Wolf Mueller⁸, Oliver Grauer⁹, Mirjam Renovanz¹⁰, Ghazaleh Tabatabai¹⁰, Sied Kebir¹¹, Martin Glas¹¹, Sotirios Bisdas¹², Peter Hambsch¹³, Clemens Seidel¹³, Michael Hölzel², Ulrich Herrlinger⁴





treatment course of patient A-001 with

pretherapeutic tumor tissue of the same confirms distinct

